

Dynamical Friction and Cooling Flows in Galaxy Clusters

Woong-Tae Kim

Astronomy Program, SEES, Seoul National University, Seoul 151-742, Korea

*Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138,
USA*

wkim@astro.snu.ac.kr

Amr A. El-Zant

CITA, University of Toronto, Ontario M5S 3H8, Canada

and

Marc Kamionkowski

Mail Code 130-33, California Institute of Technology, Pasadena, CA 91125, USA

ABSTRACT

We investigate a model of galaxy clusters in which the hot intracluster gas is efficiently heated by dynamical friction (DF) of galaxies. We allow for both subsonic and supersonic motions of galaxies and use the gravitational drag formula in a gaseous medium presented by Ostriker (1999). The energy lost by the galaxies is either redistributed locally or into a Gaussian centered on the galaxy. We find that the condition of hydrostatic equilibrium and strict energy balance yields a trivial isothermal solution T_{iso} , independent of radius, or rising temperature distributions provided $T_{\text{iso}}/\gamma < T < T_{\text{iso}}$, where γ is the adiabatic index of the gas. The isothermal temperature corresponds to the usual scaling relation between the gas temperatures and the velocity dispersions of galaxies. However the minimal temperature associated with the rising solutions is $\sim \frac{1}{2}T_{\text{vir}}$, larger than that inferred from observations, the radial distribution of galaxy masses notwithstanding. Heating by supersonically moving galaxies cannot suppress thermal instability, although it can lengthen the growth time up to the level comparable to the ages of clusters when Mach number of galaxies is less than about two. We show using numerical hydrodynamic simulations that DF-induced heating is generally unable to produce stable equilibrium cores by evolving arbitrary non-equilibrium clusters, although it can lengthen the cooling time. We conclude that

DF-induced heating *alone* is an unlikely solution to the cooling flow problem, although it can still be an important heat supplier, considerably delaying cooling catastrophe. We discuss other potential consequences of DF of galaxies in galaxy clusters.

Subject headings: galaxies: clusters, general — galaxies: interactions — galaxies: kinematics and dynamics — cooling flows — X-rays: galaxies — instability

1. Introduction

Galaxy clusters are the largest gravitationally bound structures in the universe. They typically contain several hundreds to thousands of galaxies orbiting in a gravitational potential well formed primarily by dark matter (e.g., Bahcall 1977). They are also filled with hot gas with $T \sim 2 - 10$ keV that loses thermal energy prolifically by emitting X-rays. In the absence of any heat sources, the radiative cooling in the cores of rich clusters would result in a cooling flow in which gas settles in the gravitational potential and drops out as cold condensations (e.g., Fabian 1994). However, recent high resolution *Chandra* and *XMM-Newton* observations have not shown the expected signature of gas cooling below one-third of the virial temperature (Blanton et al. 2001; Peterson et al. 2001, 2003; Tamura et al. 2001; Johnstone et al. 2002; Allen et al. 2001). This strongly suggests that there must be some source of heat that manages to balance the radiative cooling and thus prevents the mass dropout at the central regions of galaxy clusters. Most theoretical models proposed so far for cluster heating fall into the following two groups: (1) energy injection via radiations, jets, outflows, bubbles, or sound waves from a central active galactic nucleus (Ciotti & Ostriker 2001; Churazov et al. 2002; Kaiser & Binney 2003; Ruszkowski et al. 2004; Roychowdhury et al. 2004; Vecchia et al. 2004 and references therein); (2) diffusive transport of heat from the outer regions of the cluster to the center via conduction (Tucker & Rosner 1983; Bregman & David 1988; Narayan & Medvedev 2001; Voigt et al. 2002; Zakamska & Narayan 2003; Kim & Narayan 2003a) or turbulent mixing (Cho et al. 2003; Kim & Narayan 2003b; Voigt & Fabian 2004; Dennis & Chandran 2005).

Although less well studied than those mentioned above, there are clearly other energy sources in the cluster environments. These include kinetic energy in the orbital motions of galaxies (Miller 1986; Just et al. 1990), gravitational potential energy of the gas (Markevitch et al. 2001; Fabian 2003), feedback from intracluster supernovae (Domainko et al. 2004), dissipation of turbulent energy driven by infall of subclusters (Fujita et al. 2004), etc. Amount of available energy in each source is comparable to or larger than thermal energy of the intracluster medium (ICM), so that radiative cooling would be easily offset if there exist

physical mechanisms that can tap the energy from the sources and convert it into thermal energy in the ICM.

One such process is dynamical friction (DF) which occurs when a galaxy moving around a cluster center interacts with its gravitationally induced wake in the ICM (e.g., Dokuchaev 1964; Hunt 1971; Ruderman & Spiegel 1971; Rephaeli & Salpeter 1980). Because of the gravitational drag, the galaxy loses some of its kinetic energy. Turbulence in the ICM generated by the superposition of the wakes of many galaxies or by other processes will absorb the lost kinetic energy and, in the presence of tangled magnetic fields and/or viscosity, turns it into heat at the dissipation scales (e.g., Deiss & Just 1996). Miller (1986) and Just et al. (1990) independently estimated the heating rates by DF of galaxies and found that the DF-induced heating can fairly well balance radiative cooling of a rich cluster, provided the mass-to-light ratio is about 20. Using updated data for the properties of galaxies and gas in the Perseus cluster and employing Monte-Carlo approaches, El-Zant, Kim, & Kamionkowski (2004, hereafter EKK04) very recently showed that the total power generated by DF within the cooling radius is comparable to radiative loss from that region if the average mass-to-light ratio of galaxies is about 10. They also noted that this mechanism is self-regulating in the sense that gas would not be overheated since the DF of galaxies becomes ineffective when it attains high enough temperature for which galaxy motions become subsonic (see also Miller 1986).

While the results of the aforementioned work suggest that the *total* power supplied by the motions of member galaxies is comparable to X-ray luminosity of a typical rich cluster, it is still uncertain whether it can balance the radiative cooling *locally* as well. If the DF of galaxy motions is to serve as a primary heat supplier in clusters in equilibrium, observed density and temperature profiles of clusters should be a natural consequence of the local heat balance between radiative cooling and DF-induced heating. Since optically-thin, X-ray emitting gas is prone to thermal instability (e.g., Fabian 1994), it is also an interesting question whether the DF-induced heating is able to suppress thermal instability completely or at least lengthen its growth time to the level comparable to the ages of clusters; even an equilibrium cluster would otherwise be subject to a mass dropout at its center (Kim & Narayan 2003a).

In this paper, we take one step further from Miller (1986) and EKK04 to investigate equilibrium cluster models in which DF-induced heating is a main heating mechanism. We will first construct density and temperature distributions of the gas in clusters that are in strict hydrostatic and thermal equilibrium and compare them with observed profiles. We note that recent X-ray data from *BeppoSAX*, *Chandra*, and *XMM-Newton* observations indicate that gas temperature in rich clusters rapidly increases with radius for $r \lesssim (0.05 - 0.08)r_{180}$,

remains roughly isothermal up to $r \sim 0.2r_{180}$, and then gradually declines farther out (Ettori et al. 2000; De Grandi & Molendi 2002; Piffaretti et al. 2005; Vikhlinin et al. 2005, see also Markevitch et al. 1998 for *ASCA* results). Here, r_{180} is a virial radius the interior of which has the mean density of gas equal to 180 times the critical density. While declining temperature profiles at large radii may provide important clues to cosmological structure formation, they appear not to be directly relevant to the cooling flow problem. This is because gas density in clusters decreases as or faster than $\sim r^{-2}$ at the outer parts, so that the radiative cooling time in the $r \gtrsim 0.2r_{180}$ regions are much longer than the Hubble time. Indeed, the results of numerical simulations for cluster formation show that the shapes of declining temperature profiles are essentially independent of the presence of radiative cooling and/or supernova feedback (e.g., Loken et al. 2002; Motl et al. 2004). For this reason, when we compare our model results with observations, we pay attention to density and temperature profiles only in cooling regions with $r \lesssim 0.2r_{180}$ (typically $\sim 0.5 - 1$ Mpc) beyond which the effect of radiative cooling is not serious and thus heating mechanisms are not required.

The remainder of this paper is organized as follows: In §2 we evaluate the total heating rate resulting from the DF of galaxies using the formula given by Ostriker (1999) for gravitational drag force in a gaseous medium. While Miller (1986) and EKK04 approximated galaxy motions as being highly supersonic, we allow for both subsonic and supersonic motions of galaxies. In §3 we calculate equilibrium density and temperature profiles of galaxy clusters by assuming that the DF-induced heating is deposited at the locations of galaxies. Effects of distributed heating of DF and the radial mass variation of galaxies are discussed in §4. In §5 we analyze local thermal stability of the gas while in §6 we present the results of numerical hydrodynamic simulations that investigate the evolution of initially out of equilibrium configurations. Finally, in §7 we conclude this work and discuss other potential consequences of DF of galaxies in clusters.

2. Total Heating Rate

Galaxies orbiting around the center of a galaxy cluster gravitationally induce wakes in the ICM. The gravitational interaction between the galaxies and wakes causes the former to lose their orbital kinetic energy and converts it into thermal energy of the ICM via either compressional heating (including shock) or turbulent dissipation of the gas kinetic energy in the wakes. Although the notion that DF of galaxies can generate heat in the hot ICM has been well recognized and widely studied, most of previous work on this subject assumed galaxies in highly supersonic motions, estimated only total heating rate, and compared it with total radiative loss rate in clusters (Ruderman & Spiegel 1971; Rephaeli & Salpeter

1980; Miller 1986; EKK04). However, the motions of galaxies are only slightly supersonic, with an average Mach number of about 1.5 (e.g., Sarazin 1988; Faltenbacher et al. 2004), and even become subsonic at the outer parts of clusters. In this paper we consider both subsonic and supersonic galaxy motions and adopt the general formula of Ostriker (1999) for the DF force in a gaseous medium.

Following Ostriker (1999), we consider a galaxy of mass M_g moving at velocity $\mathbf{v}$ through a uniform medium with density ρ and sound speed c_s . The dynamical-friction force that the galaxy experiences is given by

$$\mathbf{F}_{\text{DF}} = -\frac{4\pi\rho(GM_g)^2 I}{v^3} \mathbf{v}, \quad (1)$$

where the efficiency factor I is defined by

$$I \equiv \begin{cases} \frac{1}{2} \ln \left(\frac{1+\mathcal{M}}{1-\mathcal{M}} \right) - \mathcal{M}, & \mathcal{M} < 1 \\ \frac{1}{2} \ln (1 - \mathcal{M}^{-2}) + \ln (vt/r_{\min}), & \mathcal{M} > 1, \end{cases} \quad (2)$$

with $\mathcal{M} \equiv v/c_s$ being the Mach number of the galaxy motion and $r_{\min}$ the effective size of a galaxy (Ostriker 1999). Note that for $\mathcal{M} \gg 1$, equations (1) and (2) with $vt = r_{\max}$ corresponding to the maximum system size, become identical to the Chandrasekhar (1943) formula for the drag force due to collisionless particles. Although equation (1) is valid for a perturber moving on rectilinear trajectory through a uniform-density medium, numerical simulations show that it is a good approximation even in a radially-stratified, spherical system (Sánchez-Salcedo & Brandenburg 2001).

We assume that the orbital velocities of N_g galaxies that contribute to heating of the gas are described by the Maxwellian distribution,

$$f(v) = \frac{4\pi N_g}{(2\pi\sigma_r^2)^{3/2}} v^2 e^{-v^2/(2\sigma_r^2)}, \quad (3)$$

with a one-dimensional velocity dispersion σ_r . The total heating rate due to the DF is then given by

$$\langle \dot{E} \rangle = N_g \langle -\mathbf{F}_{\text{DF}} \cdot \mathbf{v} \rangle = \frac{4\pi\rho(GM_g)^2 N_g}{c_s} \left\langle \frac{I}{\mathcal{M}} \right\rangle, \quad (4)$$

where the angular brackets denote an average over equation (3).

Using equations (2) and (3), one can evaluate $\langle I/\mathcal{M} \rangle$ numerically. Figure 1 plots $\langle I/\mathcal{M} \rangle$ as functions of $m \equiv \sigma_r/c_s$ for some values of $\ln(vt/r_{\min})$. For supersonic cases, the density perturbations in the wake are highly asymmetric and the regions influenced by the perturber shrink as m increases. This results in a smaller heating rate for larger m . In this case, one

can show $\langle I/\mathcal{M} \rangle \approx (2/\pi)^{1/2} m^{-1} \exp(-0.5/m^2) \ln(vt/r_{\min})$ for $m \gg 1$. For a subsonically moving perturber, on the other hand, the perturbed density in the front and back sides of the perturber becomes more or less symmetric, producing a smaller heating rate with decreasing m . In the limit of vanishingly small m , $\langle I/\mathcal{M} \rangle \rightarrow m^2$. In general, $\langle I/\mathcal{M} \rangle$ peaks at $m = 1$, as Figure 1 indicates.

Taking $vt = 1$ Mpc from a typical size of clusters and $r_{\min} = 10$ kpc as the galaxy size, $\ln(vt/r_{\min}) \approx 4.6$. Since m usually varies from 0.8 to 3, $\langle I/\mathcal{M} \rangle \approx 2$. Therefore, we have the total heating rate

$$\langle \dot{E} \rangle \approx 4 \times 10^{44} \text{ ergs s}^{-1} \left(\frac{N_g}{500} \right) \left(\frac{M_g}{10^{11} M_\odot} \right)^2 \left(\frac{n_e}{0.01 \text{ cm}^{-3}} \right) \left(\frac{T}{5 \text{ keV}} \right)^{-1/2} \left(\frac{\langle I/\mathcal{M} \rangle}{2} \right), \quad (5)$$

where n_e and T denote the electron number density and gas temperature, respectively, and we adopt the solar abundance. Notice that the rate of gas heating given in equation (5) is similar to typical X-ray luminosity of rich clusters (e.g., Sarazin 1988; Rosati, Borgani, & Norman 2002), implying that there is a sufficient amount of energy available in the orbital motions of galaxies. This is essentially what led EKK04 to conclude that the dynamical-friction coupling between cluster galaxies and gas can provide thermal energy enough to compensate for radiative loss. However, it is still questionable whether DF of galaxies heats the gas in a right manner. That is, are the observed density and temperature profiles of the ICM a direct consequence of energy balance between heating by DF and radiative cooling? Is the intracluster gas heated by DF thermally stable? In what follows, we shall study models of clusters with DF-induced heating in detail by assuming hydrostatic equilibrium and thermal energy balance.

3. Model With Localized Heating

DF of galaxies is mediated by gravity which is a long-range force, so that heating of gas due to a single galaxy is likely to be well distributed throughout its wake. The functional form of heat distribution is unknown and its derivation may require numerical simulations, which are beyond the scope of the present work. In this section, we instead make a simplifying assumption that heat is deposited *locally* at the galaxy position. The effects of heat distribution will be discussed in §4.

3.1. Local Heating Function

Consider a spherically symmetric distribution of galaxies, with the number density given by $n_g(r)$ at radius r . The local heating rate per unit volume due to DF is given by

$$\langle \dot{e}(r) \rangle = \frac{4\pi\rho(GM_g)^2\langle I/\mathcal{M} \rangle}{c_s} n_g(r), \quad (6)$$

where we assume equal galaxy mass. To find $n_g(r)$, we assume that the orbits of galaxies are isotropic and isothermal. The usual Jeans equation (e.g., eq. [4-55] of Binney & Tremaine 1987) then reads

$$\sigma_r^2 \frac{d \ln n_g}{dr} = -\frac{d\Phi}{dr}, \quad (7)$$

where Φ is the gravitational potential.

Under the NFW distribution of dark matter

$$\rho_{\text{NFW}} = \frac{M_0/2\pi}{r(r+r_s)^2}, \quad (8)$$

with a scale radius r_s and a characteristic mass M_0 , the gravitational acceleration is given by

$$-\frac{d\Phi}{dr} = -\frac{2GM_0}{r_s^2} \left[\frac{\ln(1+x)}{x^2} - \frac{1}{x(1+x)} \right], \quad (9)$$

where $x \equiv r/r_s$ is the dimensionless radius (Navarro, Frenk, & White 1997; Klypin et al. 2001). Combining equations (7) and (9) together, one can easily find

$$n_g(x) = n_g(0)g(x) = N_g \frac{g(x)}{r_s^3 \int g(x') d^3x'}, \quad (10)$$

where

$$g(x) \equiv \left[\frac{(1+x)^{1/x}}{e} \right]^\eta, \quad (11)$$

with the dimensionless parameter η defined by

$$\eta \equiv \frac{2GM_0}{r_s \sigma_r^2}. \quad (12)$$

Note that η measures the ratio of the gravitational potential energy of a galaxy in a cluster to that galaxy's kinetic energy at $r = r_s$.

While the functional form of $g(x)$ looks strange, it behaves quite well near $x \sim 0$ and gives density profiles similar to the isothermal β -model of gas distributions (Makino et al. 1998) or to the King model of the observed galaxy distributions (e.g., Girardi et al. 1998).

For example, Girardi et al. (1998) found that the best-fit distribution of galaxies in A1795 is $n_g(r) = n_g(0)[1 + (r/R_c)^2]^{-1.27}$ with a core radius $R_c = 43$ pc, which is plotted as a solid line in Figure 2. On the other hand, X-ray and optical observations indicate $M_0 = 6.6 \times 10^{14} M_\odot$, $r_s = 460$ kpc, and $\sigma_r \approx 800$ km s $^{-1}$ for A1795 (Girardi et al. 1998; Ettori et al. 2002), corresponding to $\eta \approx 19$. Figure 2 plots as a dotted line the $g(x)$ curve with $\eta = 19$, which is in good agreement with the observed galaxy distribution.

We take $2r_s$ as the outer boundary of a cluster. Inserting equation (10) into equation (6), one obtains

$$\begin{aligned} \langle \dot{e}(x) \rangle &= \frac{4\pi\rho(GM_g)^2 N_g}{c_s r_s^3 \int g(x') d^3 x'} \left\langle \frac{I}{\mathcal{M}} \right\rangle g(x) \\ &= 1.7 \times 10^{-25} r_{s,460}^{-3} N_{g,500} M_{g,11}^2 n_e T_{\text{keV}}^{-1/2} \langle I/\mathcal{M} \rangle g(x) \text{ erg cm}^{-3} \text{ s}^{-1}, \end{aligned} \quad (13)$$

where $r_{s,460} = r_s/(460 \text{ kpc})$, $N_{g,500} = N_g/500$, $M_{g,11} = M_g/(10^{11} M_\odot)$, and T_{keV} is the temperature of the gas in units of keV. Equation (13) is our desired equation for the volume heating rate due to the DF of galaxies. Note that $\langle I/\mathcal{M} \rangle \approx 2$ at $m \sim 1$ and depends on temperature.

3.2. Equilibrium Model

We look for cluster density and temperature distributions in which the hot gas satisfies both hydrostatic equilibrium and energy balance between radiative cooling and DF-induced heating. For thermal bremsstrahlung, the rate of energy loss per unit volume is given by

$$j = 7.2 \times 10^{-24} n_e^2 T_{\text{keV}}^{1/2} \text{ erg cm}^{-3} \text{ s}^{-1}, \quad (14)$$

(Rybicki & Lightman 1979; Zakamska & Narayan 2003). From equations (13) and (14), the condition of thermal energy balance yields

$$n_e T_{\text{keV}} = 2.4 \times 10^{-2} r_{s,460}^{-3} N_{g,500} M_{g,11}^2 \langle I/\mathcal{M} \rangle g(x). \quad (15)$$

Neglecting the weak temperature dependence of $\langle I/\mathcal{M} \rangle$ and assuming that M_g does not vary with radius, equation (15) states that the gas pressure in equilibrium should trace the distribution of galaxies.

Hydrostatic equilibrium of the gas requires

$$\frac{c_0^2}{n_e} \frac{d(n_e T_{\text{keV}})}{dr} = -\frac{d\Phi}{dr}, \quad (16)$$

where $c_0 = 390 \text{ km s}^{-1}$ is an *isothermal* sound speed at $T_{\text{keV}} = 1$. Substituting equations (9) and (15) into equation (16) and using equation (11), we obtain

$$\frac{d}{dx} \ln \langle I/\mathcal{M} \rangle = -\eta \left(\frac{\sigma_r^2}{c_0^2 T_{\text{keV}}} - 1 \right) \left[\frac{\ln(1+x)}{x^2} - \frac{1}{x(1+x)} \right]. \quad (17)$$

Since $\langle I/\mathcal{M} \rangle$ depends on T_{keV} , equation (17) can be integrated as an initial value problem to yield $T_{\text{keV}}(x)$. The equilibrium profile for $n_e(x)$ can then be found from equation (15). Since $n_e(x)$ depends rather sensitively on less-well known quantities such as N_g and M_g^2 and is easily influenced by the mass profile of galaxies, we concentrate on $T_{\text{keV}}(x)$ that is independent of the properties of galaxies other than σ_r .

Equation (17) has a trivial isothermal solution

$$T_{\text{iso}} = 6.5 \left(\frac{\sigma_r}{10^3 \text{ km s}^{-1}} \right)^2 \text{ keV}, \quad (18)$$

independent of the radius. The corresponding density profile is

$$n_e = 7.4 \times 10^{-3} r_{s,460}^{-3} \sigma_{r,3}^{-2} N_{g,500} M_{g,11}^2 g(x), \quad (19)$$

where $\sigma_{r,3} = \sigma_r / (10^3 \text{ km s}^{-1})$, indicating that the gas density follows the galaxy distribution exactly. Considering uncertainties in the values of $r_{s,460}$, $\sigma_{v,1000}$, $N_{g,500}$, and $M_{g,11}$,¹ equation (19) gives $n_e \sim 0.01 - 0.2 \text{ cm}^{-3}$ at the cluster centers, which is not much different from observed values. Note that the Mach number corresponding to T_{iso} is $m = \gamma^{-1/2}$. For later purposes, we define the transonic temperature

$$T_{\text{sonic}} \equiv \frac{1}{\gamma} T_{\text{iso}}, \quad (20)$$

corresponding to unity Mach number of galaxy motions.

The isothermal solution (18) is consistent with what has long been known as the scaling relation between the gas temperature and the velocity dispersion of galaxies for many clusters (e.g., Mushotzky 1984; Edge & Stewart 1991; Lubin & Bahcall 1993; Wu et al. 1999, see also Sarazin 1988; Rosati, Borgani, & Norman 2002 for review). It explains why the values of the β ($\equiv \mu m_p \sigma_r^2 / kT$) parameter in the β models for gas distributions are close to unity (Lubin & Bahcall 1993; Bahcall & Lubin 1994; Girardi et al. 1996b, 1998). Physically, the scaling relation implies that the plasma and the galaxies are well relaxed under the common gravitational potential. It is uncertain whether the observed scaling law results truly from

¹For example, the average mass of galaxies would increase up to $\sim 5 \times 10^{11} M_\odot$ if the contribution from dark halos is taken into account (e.g., Zentner & Bullock 2003).

the DF coupling between the ICM and galaxies, but equation (18) suggests that the latter certainly makes the former tighter.

Although gas in the regions surrounding cooling cores of clusters has nearly constant temperatures close to the virial values, recent X-ray observations exhibit positive radial gradients of gas temperatures at the central ~ 300 kpc regions (Peterson et al. 2001, 2003), while showing declining temperature distributions at far outer regions (Markevitch et al. 1998; De Grandi & Molendi 2002; Piffaretti et al. 2005; Vikhlinin et al. 2005). Thus, the isothermal solution cannot describe the observed temperature distributions for the whole range of radii. We are particularly interested in the regions within and adjacent to the cooling radius in which gas density is high enough to experience significant radiative cooling. To check whether the general solutions of equation (17) produce temperature and density distributions similar to the observed in such regions, we solve it numerically. We adopt $\eta = 19$, $r_s = 460$ kpc, $N_g = 500$, $M_g = 10^{11} M_\odot$, and $\sigma_r = 10^3$ km s $^{-1}$ corresponding to $T_{\text{iso}} = 6.5$ keV. We choose a value for the central temperature $T(0)$ and then integrate equation (17) from $r = 0$ to $2r_s$. The resulting temperature and density profiles for a few selected values of $T(0)$ are plotted in Figure 3.

The spatial behavior of the solutions depends critically on $T(0)$. When $T(0) > T_{\text{sonic}}$, galaxy motions are subsonic and $d\langle I/\mathcal{M} \rangle/dm > 0$ from Figure 1. If $T(0) > T_{\text{iso}}$ (or $m < \gamma^{-1/2}$), equation (17) gives $d\langle I/\mathcal{M} \rangle/dr = -(m/2T)(d\langle I/\mathcal{M} \rangle/dm)(dT/dr) > 0$ for $T(0) > T_{\text{iso}}$, implying $dT/dr < 0$. As temperature monotonically decreases with radius, m increases until it reaches the isothermal value $\gamma^{-1/2}$ where $T = T_{\text{iso}}$. One can similarly show that $dT/dr > 0$ for $T_{\text{iso}} > T(0) > T_{\text{sonic}}$, and thus temperature slowly increases toward T_{iso} . As long as $T(0) > T_{\text{sonic}}$, the solutions asymptote to the isothermal ones (eqs. [18] and [19]) as the radius increases. On the other hand, $dT/dr < 0$ when $T(0) < T_{\text{sonic}}$ (or $m > 1$). Since m increases with decreasing T and since $d\langle I/\mathcal{M} \rangle/dr$ does not change its sign for $m > 1$, $dT/dr < 0$ is satisfied for all radii. The decrease of temperature is much faster than that of $g(x)$, resulting in a unrealistic distribution of electron number density that increases with radius.

As Figure 3 shows, the local heat balance with DF-induced heating yields rising temperature profiles, if and only if $T_{\text{sonic}} < T < T_{\text{iso}}$. While this is a tantalizing result considering the central depression of temperatures seen in cooling-flow clusters, the required range of temperatures is very narrow. If we identify T_{iso} with the virial temperature, the central temperatures in our equilibrium models must be larger than $T_{\text{sonic}} = 0.6T_{\text{iso}}$ (for $\gamma = 5/3$), which is about twice larger than the observed values of $T(0) \sim (0.3 - 0.4)T_{\text{iso}}$ (e.g., Peterson et al. 2001, 2003; Allen et al. 2001). By having too stringent upper and lower limits of temperatures, therefore, heating by DF of galaxies alone is unlikely to explain observed tem-

perature and density distributions of galaxy clusters. We note however that the tight range of temperatures for the rising temperature profiles may be caused by the basic assumptions of this section, namely that DF-induced heating of the ICM is all localized at the positions of galaxies, and that all galaxies have equal mass. We relax these assumptions in the next section.

4. Effects of Distributed Heating

In the presence of viscosity and/or turbulent magnetic fields in the ICM, the kinetic energy lost by a galaxy via DF will eventually be converted into heat, rather than bulk motions distributed onto the gravitational wakes of the galaxy. Deiss & Just (1996) used a quasi-linear fluctuation theory to derive the spatial structure (in Fourier space) of mechanical heating in a turbulent medium self-consistently driven by DF of many galaxies. To our knowledge, there is no published work that study spatial heat distribution (in real space) caused by DF. If ICM heating by DF of galaxies occurs through turbulent dissipation of gas kinetic energy, finding the functional form of the heat distribution would not be viable unless the characteristics of ICM turbulence and related processes are prescribed (Deiss, Just, & Kegel 1990). Instead of attempting to derive a realistic heat distribution, in this work we simply adopt a Gaussian function to parametrize the spatial extent to which heat is distributed². Our aim in this section is to examine the effects of heat distribution on equilibrium structures in comparison with the localized heating models. Since masses of galaxies inside a cluster are likely to vary with the radius within the cluster, we also allow for spatially varying galaxy masses.

Assuming that heat distribution follows a Gaussian profile centered at the location of the galaxy, we write the heating rate per unit volume as

$$\langle \dot{e}(r) \rangle = \frac{4\pi\rho G^2 \langle I/\mathcal{M} \rangle}{c_s} h(r), \quad (21)$$

where

$$h(r) \equiv \frac{1}{\pi^{1/2} l_h} \int n_g(r') M_g(r')^2 e^{-(r-r')^2/l_h^2} dr', \quad (22)$$

is a convolution of $n_g M_g^2$ with the Gaussian function with scale length l_h . In writing equations (21) and (22), we assume that density and temperature of the gas do not change significantly

²We also tried with logarithmic heat distribution functions used in EKK04 and found that results are similar to those with Gaussian functions presented in this section.

inside the wake of a galaxy, which is acceptable within the framework of equation (1). The condition of hydrostatic equilibrium and thermal energy balance is then reduced to

$$\frac{d}{dx} \ln \langle I/\mathcal{M} \rangle = -\eta \frac{\sigma_r^2}{c_0^2 T_{\text{keV}}} \left[\frac{\ln(1+x)}{x^2} - \frac{1}{x(1+x)} \right] - \frac{d \ln h(x)}{dx}. \quad (23)$$

It can be easily verified that equation (23) recovers equation (17) in the limit of $l_h \rightarrow 0$.

Figure 4 plots the solutions of equation (23) for the case of constant M_g . More widely distributed heating is equivalent to localized heating with a flatter distribution of galaxy number density, causing the $|d \ln h/dx|$ term in equation (23) to be smaller. Consequently, an equilibrium cluster with a larger value of l_h tends to make $\langle I/\mathcal{M} \rangle$ decreases faster with increasing r . This implies that for $T(0) < T_{\text{sonic}}$ ($T(0) > T_{\text{sonic}}$), the temperature in models with non-zero l_h decreases (increases) more rapidly than in the $l_h = 0$ counterpart, although the difference between models with $l_h/r_s = 0.1$ and 1.0 is negligible for small r . Note that even when $T(0) > T_{\text{iso}}$, for which localized-heating models have declining temperature profiles, equilibrium temperature in distributed-heating models increases in the regions with $r < l_h$ and eventually converges to T_{iso} at $r \gtrsim (2-3)l_h$. We see that distributed heating does not change the condition $T_{\text{sonic}} < T < T_{\text{iso}}$ for the existence of rising temperature distributions. The distributed-heating models do not work better than the localized-heating model in terms of producing temperature distributions similar to observations; it in fact aggravates the situation by making all of the gas in clusters nearly isothermal except only in the central ~ 1 kpc regions.

Another factor that may affect the equilibrium structure is the radial mass variation of cluster galaxies. Although galaxies located near the central parts of clusters tend to have a large fraction of luminous matter, (e.g., Biviano et al. 2002; Mercurio et al. 2003), the strong tidal stripping of dark-matter halos is likely to cause the total (luminous + dark) masses of individual galaxies to be smaller toward the cluster center (Zentner & Bullock 2003; EKK04; Nagai & Kravtsov 2005). Motivated by this consideration, we adopt several algebraic forms of $M_g(r)$ that either monotonically decrease or increase by about a factor of 20 or less from the cluster center to $r = 2r_s$, and then solve equation (23). The resulting temperature profiles are almost identical to those of the constant-mass cases presented in Figure 4. Because the distribution of galaxy masses appears logarithmically in equation (23), it does not considerably affect temperature structures, although it dramatically changes electron number-density profiles.

5. Thermal Stability

We now discuss thermal stability of a system in which radiative cooling is locally balanced by DF-mediated heating. For spatially localized heating, the net loss function $\rho\mathcal{L}$ is given by

$$\rho\mathcal{L} = j - \langle \dot{e} \rangle = \alpha \rho^2 T^{1/2} - \beta \rho T^{-1/2+p}, \quad (24)$$

where the power index p accounts for the local temperature dependence of $\langle I/\mathcal{M} \rangle$ in a piecewise manner, and the positive constants α and β contain all information on the cluster properties other than gas density and temperature. Figure 5 plots $p = d \ln \langle I/\mathcal{M} \rangle / d \ln T$ as a function of m , which gives $p \approx 0$ at around the transonic temperature ($m \approx 1$), while $p \approx 0.3 - 0.5$ for supersonic temperatures ($m > 1$). The curves asymptote to -1 for $m \ll 1$ and to 0.5 for $m \gg 1$, as the asymptotic formulae given in §2 suggest. Note that p is insensitive to the choice of $\ln(vt/r_{\min})$ as long as $m > 0.8$.

Thermal stability of a system can be easily checked by using the generalized Field criterion which states that the system is thermally unstable to isobaric perturbations if

$$\left. \frac{\partial(\mathcal{L}/T)}{\partial T} \right|_P = \frac{1}{\rho T} \left[\left. \frac{\partial(\rho\mathcal{L})}{\partial T} \right|_\rho - \frac{\rho}{T} \left. \frac{\partial(\rho\mathcal{L})}{\partial \rho} \right|_T \right] = -p \frac{\alpha \rho}{T^{3/2}} < 0, \quad (25)$$

where the second equality assumes $\rho\mathcal{L} = 0$ corresponding to thermal equilibrium (Field 1965; Balbus 1986). Since $p > 0$ for $m > 1$ (and $p < 0$ for $m < 1$), this implies that the ICM is thermally unstable (stable) if heating is caused preferentially by galaxies moving at a supersonic (subsonic) speed. Dense inner parts of galaxy clusters, where the cooling time is less than the Hubble time, are filled with low-temperature gas such that galaxy motions are readily supersonic, implying that DF-induced heating is unable to quench thermal instability in those cooling cores. Whether thermal instability has important dynamical consequences on cluster evolution can be judged by considering its growth time which amounts to

$$t_{\text{grow}} = -\frac{\gamma}{\gamma - 1} \frac{P}{\rho T^2} \left. \frac{\partial(\mathcal{L}/T)}{\partial T} \right|_P^{-1} = 0.96 p^{-1} \text{ Gyr} \left(\frac{n_e}{0.05 \text{ cm}^{-3}} \right)^{-1} \left(\frac{k_B T}{2 \text{ keV}} \right)^{1/2}, \quad (26)$$

(cf. Kim & Narayan 2003a). Even if cooling cores are thermally unstable, therefore, the virulence of thermal instability may or may not manifest itself during the lifetime of clusters depending on the value of p . For example, for a cluster with $\sigma_r = 10^3 \text{ km s}^{-1}$ and $T = 2 \text{ keV}$ at the central regions, $m = 1.4$ and $p = 0.23$ from Figure 5, giving $t_{\text{grow}} = 4.2 \text{ Gyr}$. This is almost comparable to the ages of clusters since the last major merger event ($\sim 7 \text{ Gyr}$ for massive clusters; Kitayama & Suto 1996), suggesting that thermal instability may be dynamically unimportant for practical purposes.

The inability of DF-induced heating to suppress thermal instability seems not to be a fatal problem as long as the following two conditions are met: i) cooling cores are in thermal equilibrium, a basic premise of the stability analysis; and ii) galaxies near the central parts move at near-transonic speeds ($m \lesssim 2$), so as to have a sufficiently small value of p . While the second condition appears to be easily satisfied in clusters, the results of the preceding section suggests that the first condition may not be so if DF-induced heating is to balance radiative cooling for the *whole* range of radii. Since the cooling time outside the cooling radius is longer than the Hubble time, one may argue that material beyond the cooling radius does not need any heating source and that it is sufficient for DF of galaxies to supply heat to gas only in cooling cores. We address this possibility in the next section.

As a final remark of this section, we compare our results with previous work. Formally, equation (25) implies that gas subject to DF-induced heating has a critical temperature above (below) which it becomes thermally stable (unstable) and that this critical temperature corresponds exactly to unity Mach number of galaxy motions. This result is quite different from those of previous work which made some approximations on the heating function. For instance, Miller (1986) argued that gas with drag heating is always thermally unstable and rapidly turns into a multi-phase medium, a consequence of his neglecting the temperature dependence of the heating function (corresponding to $p = 1/2$ in eq. [24]). Although Bregman & David (1989) found a similar critical gas temperature, they considered only the supersonic regime and used an approximate heating function from the shock jump conditions, which makes their critical temperature smaller than ours by a factor of about three. Our result, based on the general formula of Ostriker (1999) for the drag force applicable for both supersonic and subsonic cases, represents the situation better.

6. Time Dependent Approach

We have shown in the preceding sections that an equilibrium cluster subject to DF-mediated heating has a temperature profile that either decreases (for $T > T_{\text{iso}}$ or $T < T_{\text{sonic}}$) or increases (for $T_{\text{sonic}} < T < T_{\text{iso}}$) with radius. While the radially increasing temperature profile is attractive, the requirement that thermal energy is balanced locally in the entire regions out to $2r_s$ causes the central gas temperature to be no less than 0.6 times the virial temperature, about a factor two larger than observations. Since radiative cooling is important only in the dense central parts inside the cooling radius, however, it would be sufficient for DF of galaxies to provide heat only in such a cooling core instead of the entire regions. In addition, thermal instability would not be an issue provided that gas is in thermal equilibrium and that galaxy motions are near transonic (§5). Therefore, it may be possible

for a cluster to start from an arbitrary non-equilibrium state and evolve slowly under both radiative cooling and DF-mediated heating, ending up with an equilibrium core in which thermal instability has a very long growth time. The resulting temperature profile, albeit consistent with observations, need not represent an equilibrium for the whole range of radii.

To test this idea, we have run a number of one-dimensional hydrodynamic simulations in spherical polar coordinates. We set up a logarithmically spaced radial grid, with 400 zones, from 1 kpc to 1 Mpc, and explicitly implement the net cooling-heating function described by equation (24). We also include the effect of gravitational potential due to *passive* dark matter using equation (9) with $M_0 = 6.6 \times 10^{14} M_\odot$ and $r_s = 460$ kpc; the DF-coupling between gas and the dark matter and back reaction on the latter component are not taken into account. As initial conditions, we consider spherically symmetric clusters and take $n_e = n_e(0)g(x) \text{ cm}^{-3}$ with the central electron density $n_e(0) = 0.002, 0.005, 0.01, 0.02 \text{ cm}^{-3}$, and fix $T = 6.5 \text{ keV}$ independent of the radius for all models. We adopt $r_s = 460$ kpc, $N_g = 500$, and $M_g = 10^{11} M_\odot$, and consider both localized heating and distributed heating with $l_h/r_s = 0.1$ and 1. Note that the corresponding equilibrium density profile has $n_e(0) = 0.0074 \text{ cm}^{-3}$ when heating is localized. We adopt the outflow boundary conditions for scalar variables (i.e., density, energy, etc.) that assign the same values in the ghost zones as in the corresponding active zones. For the radial velocity, we allow it to vary as a linear function of radius at the inner boundary, while fixing it to be zero at the outer boundary (e.g., Kim & Narayan 2003a). Using the ZEUS hydrodynamic code (Stone & Norman 1992), we solve the basic hydrodynamic equations and follow the nonlinear evolution of each model cluster. The resulting radial profiles of electron number density and temperature of model A with $n_e(0) = 0.02 \text{ cm}^{-3}$ and model B with $n_e(0) = 0.005 \text{ cm}^{-3}$, both of which assume spatially localized heating, are shown in Figure 6 for a few time epochs.

Model A, which initially has larger density than the equilibrium value everywhere, immediately develops radial mass inflows. As time evolves, the temperature drops and the density increases. Compared to pure cooling cases which fairly well maintain near isobaric conditions (e.g., Kim & Narayan 2003a), DF-induced heating is found to cause thermal pressure to increase with density as $P \propto \rho^{0.3}$. Since $j \propto T^{-2.4}$ and $\langle \dot{e} \rangle \propto T^{-1.9+p}$ with $p > 0$ for supersonic galaxy motions, however, the system is thermally unstable and cooling occurs at a much faster rate as temperature decreases. Model A experiences a cooling catastrophe in less than 4 Gyr. On the other hand, model B with initial overheating becomes hotter and less dense, which in turn tends to increase the ratio of heating rate to the cooling rate for $p \gtrsim -0.5$. As the gas temperature increases, the Mach number of galaxy motions become smaller, reducing the value of p . The cooling rate will therefore eventually exceed the heating rate and the cluster evolution will be reversed. In the case of model B, this turnaround will occur at $t = 70 \text{ Gyr}$ when $T \sim 18 \text{ keV}$ (or $m \sim 0.5$) is reached.

The evolution of models with different density and different length scales for heat distribution is qualitatively similar to those of models A and B, namely that clusters catastrophically evolve toward vanishing central temperature if cooling dominates initially, while clusters with initial excessive heating heat up steadily. This suggests that DF-induced heating does not naturally lead non-equilibrium clusters to thermally stable, equilibrium cores. As shown in §3 and §4, DF of galaxies does not explain the observed temperature distributions of clusters if the condition of thermal energy balance is imposed for all radii. Although it is enough for DF-induced heating to balance radiative cooling only in cooling cores, such cooling cores do not naturally form from non-equilibrium states. Unless the properties and galaxies and ICM are fine tuned, small departures from an equilibrium state rapidly evolve into an extreme configuration. Therefore, we conclude that DF-induced heating *alone* is not likely to account for the absence of cold gas in the centers of galaxy clusters.

Although heating by DF from galaxies does not appear to provide a complete solution to the cooling flow problem, we see that the DF-induced heating can still offset a considerable amount of radiative cooling. The isobaric cooling time is given by

$$t_{\text{cool}} = \frac{\gamma}{\gamma - 1} \left(\frac{P}{\rho \mathcal{L}} \right) = \frac{0.96}{1 - \mathcal{R}/(n_e T_{\text{keV}})} \left(\frac{n_e}{0.05 \text{ cm}^{-3}} \right)^{-1} \left(\frac{kT}{2 \text{ keV}} \right)^{1/2} \text{ Gyr}, \quad (27)$$

where $\mathcal{R} \equiv 0.024 r_{s,460}^{-3} N_{g,500} M_{g,11}^2 \langle I/\mathcal{M} \rangle$ represents the contribution of the DF-induced heating. Figure 7 plots as solid lines the cooling time as a function of n_e for $N_{g,500} M_{g,11}^2 = 1$ or as a function of $N_{g,500} M_{g,11}^2$ with varying density. The cases without any heating are compared as dotted lines. The time epochs when clusters experience cooling catastrophe in the numerical simulations are marked in Figure 7a as solid circle, triangle, cross, and open circle for no heating, the distributed heating with $l_h/r = 1.0$ and 0.1 , and the localized heating cases, respectively. The numerical results are in good agreement with the theoretical prediction. Models with distributed heating tend to have longer cooling time. When $N_{g,500} M_{g,11}^2 = 1$ and $n_e(0) = 0.05 \text{ cm}^{-3}$, Figure 7a shows that DF-induced heating lengthens the cooling time by about 13% compared to the pure cooling case. However, as Figure 7b indicates, the offset of cooling by DF-induced heating is increasingly larger as $N_{g,500} M_{g,11}^2$ increases. The increment of the cooling time could be as large as 140% when $n_e(0) \sim 0.03 - 0.1 \text{ cm}^{-3}$ and $N_{g,500} M_{g,11} \sim 3 - 7$, suggesting that thermal energy supplied by DF of galaxies is by no means non-negligible.

7. Conclusions and Discussion

Friction of galaxy motions via the gravitational interaction with their own gravitationally induced wake in the ICM has often been invoked as an efficient heating mechanism of

the gas (e.g., Miller 1986; Just et al. 1990; Deiss & Just 1996; EKK04). This idea of DF-mediated heating of the ICM is quite attractive because there is sufficient energy available in galaxy motions and the mechanism is self-regulating; it operates effectively only when the temperature of gas is in a certain range, which happens to be the typical gas temperatures in the cooling cores of galaxy clusters. In this paper, we take one step further from Miller (1986) and EKK04 to calculate equilibrium density and temperature profiles of the hot gas heated by DF of galaxies. Instead of restricting to cases where galaxy motions are all supersonic, we use the general formula derived by Ostriker (1999) for the drag force that takes allowance for both supersonic and subsonic galaxy motions in a gaseous medium. We show that the total heating rate due to the DF of galaxies in a typical rich cluster is comparable to its total X-ray luminosity, confirming the results of Miller (1986) and EKK04 (see §2).

Next, we derive the local heating function (eq. [13]) assuming that the orbits of galaxies are isotropic and isothermal under the NFW distribution of dark matter and that the kinetic energy lost by a galaxy is deposited into heat at the location of the galaxy. The condition that the gas is in hydrostatic equilibrium and maintains energy balance requires the temperature profile to be one of the following three kinds: isothermal, with $T = T_{\text{iso}}$; a decreasing profile with radius when $T < T_{\text{sonic}}$ or $T > T_{\text{iso}}$; an increasing profile when $T_{\text{sonic}} < T < T_{\text{iso}}$, where T_{iso} and T_{sonic} denote the temperatures corresponding to unity Mach number of galaxy motions with respect to the isothermal and adiabatic sound speeds, respectively (eqs. [18] and [20]). The isothermal solution is interesting because it quite well describes the observed scaling relationship among clusters properties. Although the rising temperature solution is attractive since clusters usually show a temperatures drop at the central regions, it strictly requires the central temperature to be no lower than 0.6 times the virial temperature, which is roughly twice smaller than the observed values (§3). We also consider cases in which DF-induced heating is distributed in space according to a Gaussian form, and/or the masses of cluster galaxies vary over radius, and find that the stringent limit of $T > T_{\text{sonic}} = 0.6T_{\text{iso}}$ for rising temperature distributions remains unchanged (§4).

Using the local heating function we have derived, we analyze thermal stability of the gas subject to radiative cooling and DF-mediated heating (§5). When galaxy motions are subsonic, the heating rate that decreases steeply with temperature suppresses thermal instability completely. On the other hand, supersonic galaxy motions for which the heating rate is relatively insensitive to temperature are unable to stop the growth of thermal instability. The growth time in the presence of DF-induced heating is at least twice that in the pure cooling case, and becomes progressively longer as the Mach number of galaxies decreases. When galaxies move at slightly supersonic velocities with Mach number less than 2, which is very likely at cluster centers, the growth time becomes comparable to the ages of clusters. This implies that thermal instability, even if operates, may have insignificant dynamical

consequences on cluster evolution.

Noting that regions outside the cooling radius need not be in strict thermal equilibrium, we look for a possibility using numerical hydrodynamic simulations that clusters evolve from an arbitrary non-equilibrium state and form current cooling cores in which DF-induced heating balances radiative cooling (§6). We find that clusters that were initially dominated by cooling unavoidably develop radial mass inflows and decrease their central temperatures in a runaway fashion, whereas clusters with initial overheating slowly heat up and result in radially decreasing temperature profiles. Equilibrium solutions therefore do not appear to form an attracting set for galaxy and gas configurations, suggesting that when DF from galaxies is the sole heating source, it is extremely difficult to obtain equilibrium cores by smoothly evolving non-equilibrium clusters (even if in some cases the cooling catastrophe can be deferred so as to occur on a longer timescale).

Putting together all the results of this paper we conclude that DF of galaxies *alone*, albeit an interesting heating mechanism with a lot of available energy, cannot be the lone heating agency to balance radiative cooling in rich galaxy clusters. We nonetheless note that the heating due DF of galaxies could considerably lengthen the cooling time, depending on the value of $N_g M_g^2$, and thus should not be neglected in energetics of galaxy clusters.

One of the key assumptions made in this paper is that all the kinetic energy lost by galaxies via DF is transferred to the thermal energy of the surrounding gas. The rationale behind this assumption is that the superposition of the wakes produces turbulence in the ICM and the kinetic energy injected into the ICM turbulence at saturation cascades down along the Kolmogorov-like energy spectrum, and eventually transforms into heat through viscous dissipation at small scales. Another possibility is that a large fraction of the kinetic energy of galaxies is used to merely enhance the level of the turbulence instead of being converted into heat. This may occur when the turbulence is not fully developed yet.

The characteristics of the ICM turbulence is not well known, yet observations and numerical simulations suggest an average velocity dispersion $v_{\text{turb}} \sim 200 - 400 \text{ km s}^{-1}$ on scales of $\lambda \sim 5 - 20 \text{ kpc}$ (Ricker & Sarazin 2001; Carilli & Taylor 2002; Churazov et al. 2004; Schuecker et al. 2004; Faltenbacher et al. 2004). The associated turbulent kinetic energy is $(1/2)M_{\text{gas}}v_{\text{turb}}^2 \sim 9 \times 10^{61} \text{ erg}$ for the total mass $M_{\text{gas}} \sim 10^{14} M_{\odot}$ of the ICM in a rich cluster. Let us assume that this energy is supplied solely by DF of galaxies at a rate given by equation (5). If the level of turbulence keeps increasing without dissipation, it would take about 7 Gyr for the DF of galaxies to feed the ICM turbulence to the observed level, which is almost comparable to the lifetime of the cluster. Conversely, if the turbulence is fully developed and in a steady state such that the energy injection rate by DF is equal to the rate of dissipation, it would have $v_{\text{turb}} = (2\lambda\dot{E}/M_{\text{gas}})^{1/3} \sim 60 \text{ km s}^{-1}$ for $\lambda \sim 20 \text{ kpc}$, which is

too small to explain the observations. All of these imply that the contribution from the DF of galaxies to the ICM turbulence is not considerable (see also Sánchez-Salcedo et al. 1999). If the dissipation of the (fully developed) turbulence is to provide enough heat to balance radiative cooling, as suggested by Churazov et al. (2004) and Fujita et al. (2004), the energy injection should not be entirely due to the DF of galaxies; it requires other energy sources, including jets from active galactic nuclei and mergers with smaller groups or clusters.

The assertion that the conversion of galaxy kinetic energy into thermal energy of the gas may not be so effective is supported by the results of Faltenbacher et al. (2004) who studied using numerical simulations cluster formation in Λ CDM cosmology, with the DF of galaxies included explicitly. They found that (1) the motions of galaxies at the present epoch are slightly supersonic, with an average Mach number of $\mathcal{M} \approx 1.4$; (2) gas within the virial radius has a velocity dispersion $v_{\text{turb}} \sim (0.3 - 0.5)c_s$ resulting probably from infall motions of galaxies and small groups, with small contribution from the DF of galaxies; and (3) the clusters still suffer from the cooling catastrophe. Although higher-resolution, more-controlled simulations are required to make a definitive statement, the last point of their results suggests that the system is thermally unstable or heating by DF may be quite inefficient — for, as we have seen in the previous section, systems evolved from arbitrary initial conditions do not generally tend to the equilibrium solutions, which do not therefore form an attracting set; and for some initial parameters the delay in the cooling catastrophe that the DF mechanism leads to is not significant.

We finally note that several issues of potential importance were not investigated in this paper. One of these relates to the galaxy velocity anisotropy. It is easy to show that, for isothermal distribution with constant anisotropy, it is possible to obtain equilibrium solutions with smaller central gas temperatures, more in line with observations. However these tend to have unrealistic galaxy number density distributions. The equilibrium and stability of gas configurations in more realistic anisotropic models have not been investigated. We have adopted the general formula of Ostriker (1999) for the drag force on a galaxy moving at an arbitrary speed. This is an improvement over previous studies (e.g., Miller 1986; EKK04) that considered only supersonic galaxy motions. While its explicit dependence on the Mach number enabled us to explore temperature profiles of equilibrium clusters, the formula still assumes a galaxy moving in a *straight-line* orbit through a *uniform* gaseous medium. Clearly, the ICM is radially stratified and galaxies follow curved rather than rectilinear trajectories. In a collisionless system, Just & Peñarrubia (2004) recently showed that the density inhomogeneity reduces the Coulomb logarithm of the Chandrasekhar formula by limiting the maximum impact parameter to the local scale length of density variation (see also Hashimoto et al. 2003; Spinnato et al. 2003), and induces an additional drag force in the lateral direction of the galaxy motion (see also Binney 1977). They further showed that the reduction

of the Coulomb logarithm makes the orbital energy loss 15% less effective, inhibiting the circularization of the orbit, while the additional tangential drag force has a negligible effect on the orbital evolution. Similar effects are expected to occur in a gaseous medium, yet their consequences are not known quantitatively. A related important question is how heat (or turbulent) energy is distributed in the wakes that form in the turbulent, inhomogeneous, magnetized ICM. A real assessment of the effects of DF to dynamical evolution of galaxy clusters requires answers to the above questions, which will direct our future research.

We are pleased to thank P. Goldreich, M. G. Lee, C. McKee, R. Narayan, and E. Ostriker for many helpful discussions and communications. We also thank an anonymous referee for useful comments and suggestions. W.-T. Kim was supported in part by Korea Science and Engineering Foundation (KOSEF) grant R01-2004-000-10490-0 at the SNU and in part by NSF grant AST 0307433 at the CfA. M. Kamionkowski was supported by NASA NNG05GF69G at Caltech.

REFERENCES

- Allen, S. W., Schmidt, R. W., & Fabian, A. C. 2001, MNRAS, 328, L37
- Bahcall, N. A. 1977, ARA&A, 15, 505
- Bahcall, N. A., & Lubin, L. M. 1994, ApJ, 426, 513
- Balbus, S. A. 1986, ApJ, 303, L79
- Binney, J. 1977, MNRAS, 181, 735
- Binney, J., & Tremaine, S. 1987, Galactic Dynamics (Princeton: Princeton Univ. Press)
- Biviano, A., Katgert, P., Thomas, T., & Adami, C. 2002, A&A, 387, 8
- Blanton, E. L., Sarazin, C. L., McNamara, B. R., & Wise, M. W. 2001, ApJ, 558, L15
- Bregman, J. N., & David, L. P. 1988, ApJ, 326, 639
- Bregman, J. N., & David, L. P. 1989, ApJ, 341, 49
- Carilli, C. L., Taylor, G. B. 2002, ARA&A, 40, 319
- Chandrasekhar, S. 1943, ApJ, 97, 255
- Churazov, E., Sunyaev, R., Forman, W., & Böhringer, H. 2002, MNRAS, 332, 729

- Ciotti, L., & Ostriker, J. P. 2001, *ApJ*, 551, 131
- Cho, J., Lazarian, A., Honein, A., Knaepen, B., Kassinos, S., & Moin, S. *ApJ*, 589, L77
- Churazov, E., Forman, W., Jones, C., Sunyaev, R., & Böhringer 2004, *MNRAS*, 347, 29
- De Grandi, S., & Molendi, S. 2002, *ApJ*, 567, 163
- Deiss, B. M., & Just, A. 1996, *MNRAS*, 305, 407
- Deiss, B. M., Just, A., & Kegel, W. H. 1990, *A&A*, 240, 123
- Dennis, J. T., & Chandran, B. D. G. 2005, *ApJ*, 622, 205
- Dokuchaev, V. P. 1964, *Soviet Astron.*, 8, 23
- Domainko, W., Gitti, M., Schindler, S., & Kapferer, W. 2004, *A&A*, 425, L21
- Edge, A. C., & Stewart, G. C. 1991, *MNRAS*, 252, 428
- El-Zant, A., Kim, W.-T., & Kamionkowski, M. 2004, *MNRAS*, 354, 169 (EKK04)
- Ettori, S., Bardelli, S., De Grandi, S., Molendi, S., Zamorani, G., & Zucca, E. 2000, *MNRAS*, 318, 239
- Ettori, S., Fabian, A. C., Allen, S. W., & Johnstone, R. M. 2002, *MNRAS*, 331, 635
- Fabian, A. C. 1994, *ARA&A*, 32, 277
- Fabian, A. C. 2003, *ARA&A*, 344, L27
- Faltenbacher, A., Kravtsov, A. V., Nagai, D., & Gottloeber, S. 2004, *MNRAS*, submitted; astro-ph/0408488
- Field, G. B. 1965, *ApJ*, 142, 531
- Fujita, Y., Matsumoto, T., & Wada, K. 2004, *ApJ*, 612, L9
- Girardi, M., Fadda, D., Giuricin, G., Mardirossian, F., & Mezzetti, M., 1996b, *ApJ*, 457, 61
- Girardi, M., Giuricin, G., Mardirossian, F., Mezzetti, M., & Boschin, W. 1998, *ApJ*, 505, 74
- Hashimoto, Y., Funato, Y., & Makino, J. 2003, *ApJ*, 582, 196
- Hunt, R. 1971, *MNRAS*, 154, 141
- Johnstone, R. M., Allen, S. W., Fabian, A. C., & Sanders, J. S. 2002, *MNRAS*, 336, 299

- Just, A., Deiss, B. M., Kegel, W. H., Böhringer, H., & Morfill, G. E., 1990, MNRAS, 354, 400
- Just, A., & Peñarrubia J. 2004, A&A, submitted; (astro-ph/0410740)
- Kaiser, C. R., & Binney, J. 2003, MNRAS, 338, 837
- Kim W.-T., & Narayan R. 2003, ApJ, 596, 889
- Kim W.-T., & Narayan R. 2003, ApJ, 596, L139
- Kitayama, T., & Suto, Y. 1996, ApJ, 469, 480
- Klypin, A., Kravtsov, A. V., Bullock, J. S., & Primack, J. R. 2001, ApJ, 554, 903
- Loken, C., Norman, M. L., Nelson, E., Burns, J., Bryan, G. L., & Motl, P. 2002, ApJ, 579, 571
- Lubin, L. M., & Bahcall, N. A. 1993, ApJ, 415, L17
- Makino, N., Sasaki, S., & Suto, Y. 1998, ApJ, 497, 555
- Markevitch, M., Forman, W. R., Sarazin, C. L., & Vikhlinin, A. 1998, ApJ, 503, 77
- Markevitch, M., Vikhlinin, A., & Mazzotta, P. 2001, ApJ, 562, L153
- Miller, L. 1986, MNRAS, 220, 713
- Mercurio, A., Massarotti, M., Merluzzi, P., Girardi, M., La Barbera, F., & Busarello G. 2003, A&A, 408, 57
- Motl, P. M., Burns, J. O., Loken C., Norman, M. L., & Bryan, G. 2004, ApJ, 606, 635
- Mushotzky, R. 1984, Phys. Scripta, T7, L157
- Nagai, D., & Kravtsov, A. V. 2005, ApJ, 618, 557
- Narayan, R., & Medvedev, M. V. 2001, ApJ, 562, L129
- Navarro, J. F., Frenk, C. S., & White, S. D. M. 1997, ApJ, 490, 493
- Ostriker, E. C. 1999, ApJ, 513, 252
- Peres, C. B., Fabian, A. C., Edge, A. C., Allen, S. W., Johnstone, R. M., & White, D. A. 1998, MNRAS, 298, 416

- Peterson, J. R., Paerels, F. B. S., Kaastra, J. S., Arnaud, M., Reiprich, T. H., Fabian, A. C., Mushotzky, R. F., & Jernigan, J. G., Sakelliou, I. 2001, *A&A*, 365, L104
- Peterson, J. R., Kahn, S. M., Paerels, F. B. S., Kaastra, J. S., Tamura, T., Bleeker, J. A. M., Ferrigo, C., & Jernigan, J. G. 2003, *ApJ*, 590, 207
- Piffaretti, R., Jetzer, Ph., Kaastra, J. S., & Tamura, T. 2005, *A&A*, 433, 101
- Rephaeli, Y., & Salpeter, E. E. 1980, *ApJ*, 240, 20
- Ricker, P. M., & Sarazin, C. L. 2001, *ApJ*, 561, 621
- Rosati, P., Borgani, S., & Norman, C. 2002, *ARA&A*, 40, 539
- Roychowdhury, S., Ruszkowski, M., Math, B. B., Begelman, M. C. 2004, *ApJ*, 615, 681
- Ruderman, M. A., & Spiegel, E. A. 1971, *ApJ*, 165, 1
- Ruszkowski, M., Brüggén M., & Begelman, M. C. 2004, *ApJ*, 615, 675
- Rybicki, G. B., & Lightman, A. P. 1979, *Radiative Processes in Astrophysics* (New York: Wiley)
- Sánchez-Salcedo, F. J., & Brandenburg, A. 1999, *ApJ*, 522, L35
- Sánchez-Salcedo, F. J., Brandenburg, A., & Suhkurov, A. 1999, *Ap&SS*, 263, 87
- Sánchez-Salcedo, F. J., & Brandenburg, A. 2001, *MNRAS*, 322, 67
- Sarazin, C. L. 1988, *X-ray emissions from clusters of galaxies* (Cambridge: Cambridge University Press)
- Schuecker, P., Finoguenov, A., Miniati, F., Böhringer, H., & Briel, U. G. 2004, *A&A*, 426, 387
- Spinnato, P. F., Fellhauer, M., Portegies Z., & Simon F. *MNRAS*, 344, 22
- Stone, J. M., & Norman, M. L. 1992, *ApJS*, 80, 753
- Tamura, T., Kaastra, J. S., Peterson, J. R., Paerels, F. B. S., Mittaz, J. P. D., Trudolyubov, S. P., Stewart, G., Fabian, A. C., Mushotzky, R. F., Lumb, D. H., & Ikebe, Y. 2001, *A&A*, 365, L87
- Tucker, W. H., & Rosner, R. 1983, *ApJ*, 267, 547

- Vecchia, C. D., Bower, R. G., Theuns, T., Balogh, M. L., Mazzotta, P., & Frenk, C. S. 2004, MNRAS, in press; astro-ph/0402441
- Vikhlinin, A., Markevitch, M., Murray, S. S., Jones, C., Forman, W., & Van Speybroeck, L. 2005, ApJin press; astro-ph/0412306
- Voigt, L., & Fabian, A. MNRAS, 347, 1130
- Voigt, L. M., Schmidt, R. W., Fabian, A. C., Allen, S. W., Johnstone, R. M. 2002, MNRAS, 335, L7
- Wu, X.-P., Xue, Y.-J., & Fang, L.-Z. 1999, ApJ, 524, 22
- Zakamska, N. L., & Narayan, R. 2003, ApJ, 582, 162
- Zentner, A., & Bullock, J. S. 2003, ApJ, 598, 49




Fig. 1.— Plots of $\langle I/\mathcal{M} \rangle$ as functions of $m \equiv \sigma_r/c_s$. Various curves corresponds to $\ln(vt/r_{\min}) = 4, 4.6, 6, 8, 10$ from bottom to top.




Fig. 2.— Comparison of $g(r)$ with $\eta = 19$ (*dotted*) with the observed distribution (*solid*) of galaxies in A1795. The core radius is taken to be $R_c = 43$ pc (Girardi et al. 1998).




Fig. 3.— (a) Equilibrium temperature and (b) electron density profiles of clusters in which DF-induced heating is spatially localized. The chosen parameters are $\eta = 19$, $r_s = 460$ kpc, $N_g = 500$, $M_g = 10^{11} M_\odot$ and $\sigma_r = 10^3$ km s $^{-1}$. Dotted and dashed lines in (a) respectively indicate the characteristic isothermal solution $T_{\text{iso}} = 6.5$ keV and the transonic temperature $T_{\text{sonic}} \equiv T_{\text{iso}}/\gamma = 3.9$ keV at which the Mach number of galaxy motions is unity. The density distribution corresponding to the isothermal solution is drawn as a dotted line in (b). Note that temperature increases with radius only when $T_{\text{sonic}} < T(0) < T_{\text{iso}}$.




Fig. 4.— Equilibrium (left) temperature and (right) density profiles of cluster models with distributed heating, which is modeled by a Gaussian function with scale length l_h . The chosen parameters are the same as in Figure 3. When $T(0) = 2, 4$ keV, temperature profiles with $l_h/r_s = 0.1$ and 1.0 are identical to each other. See text for discussion.




Fig. 5.— Logarithmic slope $p \equiv d \ln \langle I/\mathcal{M} \rangle / d \ln T = -0.5 d \ln \langle I/\mathcal{M} \rangle / d \ln m$ as a function of Mach number m for $\ln(vt/r_{\min}) = 4, 4.6, 6, 8, 10$ from top to bottom. The curves asymptote to -1 for $m \ll 1$ and to 0.5 for $m \gg 1$. With positive values of p , DF-induced heating by supersonically moving galaxies with $m > 1$ is unable to suppress thermal instability.




Fig. 6.— Evolution of electron number density (*top*) and temperature (*bottom*) for (*left*) cluster model A with $n_e(0) = 0.02 \text{ cm}^{-3}$ and (*right*) model B with $n_e(0) = 0.005 \text{ cm}^{-3}$. The initial temperature is 6.5 keV for both models. Model A, which initially has a larger density than the equilibrium value, inescapably experiences a cooling catastrophe, while model B with lower initial density steadily heats up.




Fig. 7.— Cooling time scale at the cluster center (a) as a function of n_e for fixed $N_{g,500} M_{g,11}^2 = 1$ and (b) as a function of $N_g M_g^2$ for $n_e = 0.1, 0.05, 0.01 \text{ cm}^{-3}$. Solid line is for the case when spatially localized, DF-induced heating is considered, while dotted line corresponds to the case without any heating. Various symbols in (a) represent simulation outcomes, with solid circle, triangle, cross, and open circle referring to no heating, distributed heating with $l_h/r_s = 1.0$ and 0.1 , and localized heating cases, respectively.